

# INF563 Topological Data Analysis — Exercise Session

## Homology Groups and Homotopy

### Exercise 1. Homology groups of some common spaces.

Propose triangulations of the following spaces, then compute their homology groups with coefficients in  $\mathbb{Z}/2\mathbb{Z}$ :

1. the circle (1-sphere)  $\mathbb{S}^1$ ,
2. the disk  $\mathbb{B}^2$ ,
3. the cylinder  $\mathbb{S}^1 \times [0, 1]$
4. the 2-sphere  $\mathbb{S}^2$ ,
5. the 3-ball  $\mathbb{B}^3$ ,
6. the Torus  $\mathbb{T}$ , obtained from the unit square by identifying opposite edges as illustrated in Figure 1.




Figure 1: Edge identifications on the unit square to get the torus  $\mathbb{T}$ .

### Exercise 2. Homology groups of the sphere $\mathbb{S}^d$ .

Compute the homology groups of the  $d$ -dimensional sphere  $\mathbb{S}^d$ , for any  $d \geq 1$ .

**Exercise 3. Brouwer's fixed point theorem.**

Brouwer proved the following fixed point theorem: “every continuous map  $f : \mathbb{B}^2 \rightarrow \mathbb{B}^2$  has a fixed point, i.e. a point  $p \in \mathbb{B}^2$  such that  $f(p) = p$ .” In particular, there is always a point at the surface of your coffee that is not moving. ☺

Here is a simple and elegant proof of this theorem, using homology theory. Let us assume that there exists a continuous map  $f : \mathbb{B}^2 \rightarrow \mathbb{B}^2$  that has no fixed point, and let us look for a contradiction:

1. For any  $p \in \mathbb{B}^2$ , let  $\phi(p)$  be the intersection of the open half-line  $]f(p), p)$  with the circle  $\mathbb{S}^1$ . Show that  $\phi : \mathbb{B}^2 \rightarrow \mathbb{S}^1$  is well-defined and continuous.
2. Show that  $\phi$  induces a surjective homomorphism  $\phi_*$  at the homology level. For this you can compose  $\phi$  with the canonical inclusion  $\iota : \mathbb{S}^1 \hookrightarrow \mathbb{B}^2$  and consider the induced homomorphism.
3. Conclude. Does the proof extend to higher dimensions?

**Exercise 4. The hairy ball theorem.**

The aim of this exercise is to prove the hairy ball theorem: “for  $d$  even, there is no non-vanishing continuous tangent vector field  $V : \mathbb{S}^d \rightarrow \mathbb{R}^d$ .” In particular, one cannot comb a 2-sphere that has no baldness, and there is always a point on Earth where there is no wind. ☺

Here is again a simple and elegant proof using homology theory. Let us assume that there exists a non-vanishing continuous tangent vector field  $V$  over the sphere  $\mathbb{S}^d$ , for some arbitrary  $d$ , and let us look for a contradiction when  $d$  is even.

1. Use  $V$  to find a homotopy between  $\text{id}_{\mathbb{S}^d}$  and the antipodal map  $x \mapsto -x$ .
2. Show that every morphism  $\phi_* : H_d(\mathbb{S}^d) \rightarrow H_d(\mathbb{S}^d)$  is of the form  $x \mapsto \alpha_{\phi_*} x$  for some fixed scalar  $\alpha_{\phi_*}$ .

The *degree* of a continuous map  $f : \mathbb{S}^d \rightarrow \mathbb{S}^d$  is defined as  $\deg(f) = \alpha_{f_*}$ . One can show that the degree of the antipodal map  $x \mapsto -x$  is  $(-1)^{d+1}$  (see Theorem 21.3 in “*Elements of Algebraic Topology*” by J. Munkres).

3. What is  $\deg(\text{id}_{\mathbb{S}^d})$ ? Conclude.

**Exercise 5. The dunce hat.**

The dunce hat is a classical example of a space that is contractible (homotopy equivalent to a point) but not collapsible (a stronger notion of contractibility, which plays no role in this exercise). It is obtained by identifying the edges of a triangle as illustrated in Figure 2 left.




Figure 2: Left: from the triangle to the dunce hat. Right: homeomorphic representation where the triangle has been replaced by a disk.

The aim of this exercise is to show that the dunce hat is contractible. Let us turn the initial triangle into a disk, as shown in Figure 2 right, which is only a matter of applying some homeomorphism. Then, let  $\phi : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  be the map induced by the edge identifications (whereby  $p$  is sent to the North pole and the rest of the circle is spanned by arc  $a$ ).

1. Show that there is a homotopy between  $\text{id}_{\mathbb{S}^1}$  and  $\phi$ , that is, a map  $f : [0, 1] \times \mathbb{S}^1 \rightarrow \mathbb{S}^1$  such that  $f(0, \cdot) = \text{id}_{\mathbb{S}^1}$  and  $f(1, \cdot) = \phi$ .

Let  $A \subseteq X$  be topological spaces. We say that the pair  $(X, A)$  has the *homotopy extension property* if, for any homotopy  $f : [0, 1] \times A \rightarrow Y$  and any map  $F_0 : X \rightarrow Y$  such that  $F_0|_A = f(0, \cdot)$ , there is an extension of  $F_0$  to a homotopy  $F : [0, 1] \times X \rightarrow Y$  such that  $F(t, \cdot)|_A = f(t, \cdot)$  for all  $t \in [0, 1]$ . It is known that any pair  $(X, A)$ , where  $X$  is a simplicial complex and  $A$  is a subcomplex of  $X$ , has the homotopy extension property.

2. Show that the pair  $(\mathbb{B}^2, \mathbb{S}^1)$  has the homotopy extension property. Deduce that there is a homotopy  $F : [0, 1] \times \mathbb{B}^2 \rightarrow \mathbb{B}^2$  such that  $F(0, \cdot) = \text{id}_{\mathbb{B}^2}$  and  $F(t, \cdot)|_{\mathbb{S}^1} = f(t, \cdot)$  for all  $t \in [0, 1]$ .
3. Show that the dunce hat and  $\mathbb{B}^2$  are homotopy equivalent and conclude.